

Application of difference sequences theory

Georgy Khantarchiev

January 30, 2013

Abstract

The results of difference sequences theory are applied to analytic function theory and Diophantine equations. As a result we have the equation which connects the n -th derivative of a function with the difference sequence for the values of this function. Also the results of difference sequences theory helps to discover some features of the whole kind of Diophantine equations. The method presented allows to find limits where Diophantine equation does not have integer solutions. The higher power of Diophantine equation the better this method works.

1 Introduction

I give the short introduction to difference sequences theory below to show the main idea. Let's have a look at the following system:

$$\begin{array}{cccccccc}
 & & & & \dots & \dots & \dots & \dots \\
 & & & \Delta^3 u_0, & & \Delta^3 u_1, & & \dots \\
 & & \Delta^2 u_0, & & \Delta^2 u_1, & & \Delta^2 u_2, & \dots \\
 & \Delta u_0, & & \Delta u_1, & & \Delta u_2, & & \Delta u_3, \dots \\
 u_0, & & u_1, & & u_2, & & u_3, & u_4, \dots
 \end{array}$$

where

$$\begin{aligned}
 \Delta u_k &= u_{k+1} - u_k, & u_k &= \Delta^0 u_k, \\
 \Delta^{n+1} u_k &= \Delta^n u_{k+1} - \Delta^n u_k & k &= 0, 1, 2, 3, \dots
 \end{aligned}$$

In accordance with the above definitions let's take a look at the following number sequences:

$$u_k = k, \quad k = 0, 1, 2, 3, \dots$$

$$\begin{array}{cccccccccccc} & 1, & & 1, & & 1, & & 1, & & 1, & & 1, & & \dots \\ 0, & & 1, & & 2, & & 3, & & 4, & & 5, & & 6, & \dots \end{array}$$

$$u_k = k^2, \quad k = 0, 1, 2, 3, \dots$$

$$\begin{array}{cccccccccccc} & & 2, & & 2, & & 2, & & 2, & & 2, & & \dots \\ & 1, & & 3, & & 5, & & 7, & & 9, & & 11, & \dots \\ 0, & & 1, & & 4, & & 9, & & 16, & & 25, & & 36, & \dots \end{array}$$

$$u_k = k^3, \quad k = 0, 1, 2, 3, \dots$$

$$\begin{array}{cccccccccccc} & & & 6, & & 6, & & 6, & & 6, & & \dots \\ & & 6, & & 12, & & 18, & & 24, & & 30, & \dots \\ & 1, & & 7, & & 19, & & 37, & & 61, & & 91, & \dots \\ 0, & & 1, & & 8, & & 27, & & 64, & & 125, & & 216, & \dots \end{array}$$

etc. any $u_k = k^n$, $n \in \mathbb{N}$. We can suppose that for the following number sequences: $u_k = k^n$ $k \in \mathbb{Z}$,

$$\Delta^n u_k = n!, \quad n = 1, 2, 3, \dots$$

Newton researched such difference sequences and proved the following theorem

Theorem 1.1. For polynomial $P_n(x) = a_0 x^n + a_1 x^{n-1} + \dots + a_n$, where $x \in \mathbb{R}$, $a_0, a_1, \dots, a_n \in \mathbb{R}$, $a_0 \neq 0$, $n \in \mathbb{N}$, k is the step of difference sequence, $k \in \mathbb{R}$, $k \neq 0$

$$\Delta^n P_n(x) = \sum_{i=0}^n \binom{n}{i} (-1)^i P_n(x - ki) = a_0 k^n n! \quad (1)$$

For example you can familiarize with difference sequences theory in [1].

2 Analytic Functions

Apply this result to analytic functions. Let $f(x)$ be a real analytic function in the interval (a, b) . In such case by Taylor's theorem it can be written

$$f(x) = f(x_0) + \frac{f'(x_0)}{1!}(x-x_0) + \dots + \frac{f^{(n)}(x_0)}{n!}(x-x_0)^n + \frac{f^{(n+1)}(\xi)}{(n+1)!}(x-x_0)^{n+1},$$

where $x, x_0, \xi \in (a, b)$ and the remainder has Lagrangian form.

For $i = 0, 1, 2, \dots, n$ and $k \in \mathbb{R}$, $k \neq 0$: $x + ik \in (a, b)$

$$f(x + ik) = f(x_0) + \frac{f'(x_0)}{1!}(x - x_0 + ik) + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0 + ik)^n + \frac{f^{(n+1)}(\xi_i)}{(n+1)!}(x - x_0 + ik)^{n+1},$$

where $\xi_i \in (x_0, x + ik)$. Then we can put

$$P'_n(x + ik) = \frac{f^{(n)}(x_0)}{n!}(x - x_0 + ik)^n + \dots + \frac{f'(x_0)}{1!}(x - x_0 + ik) + f(x_0),$$

hence

$$\begin{aligned} \sum_{i=0}^n \binom{n}{i} (-1)^{n-i} f(x + ik) &= \sum_{i=0}^n \binom{n}{i} (-1)^{n-i} P'_n(x + ik) + \\ &+ \sum_{i=0}^n \binom{n}{i} (-1)^{n-i} \frac{f^{(n+1)}(\xi_i)}{(n+1)!} (x - x_0 + ik)^{n+1}, \end{aligned}$$

by Theorem 1.1 and taking into account the sign at (1)

$$\sum_{i=0}^n \binom{n}{i} (-1)^{n-i} P'_n(x + ik) = f^{(n)}(x_0) k^n,$$

$$f^{(n)}(x_0)(-k)^n = \sum_{i=0}^n \binom{n}{i} (-1)^i f(x + ik) - \sum_{i=0}^n \binom{n}{i} (-1)^i \frac{f^{(n+1)}(\xi_i)}{(n+1)!} (x - x_0 + ik)^{n+1}.$$

This equation is the generalized form of (1) for functions. The equation shows a connection between the n -th derivative of a function in a point x_0 and the difference sequence for this function which begins in other point x .

In the case $x = x_0$ we have

$$f^{(n)}(x_0)(-k)^n = \sum_{i=0}^n \binom{n}{i} (-1)^i f(x_0 + ik) - \sum_{i=1}^n \binom{n}{i} (-1)^i \frac{f^{(n+1)}(\xi_i)}{(n+1)!} (ik)^{n+1}.$$

3 Diophantine equations

Lets apply the result of difference sequences theory to Diophantine equations. For example lets take Fermat's Last Theorem [2] which was proven by Wiles

in 1995 [3]. Fermat's Last Theorem states that no there positive integers x, y and z can satisfy the equation $x^n + y^n = z^n$ for any integer value of n greater than two.

Let $x > y$. For x^n we can write

$$n! = \sum_{i=0}^n \binom{n}{i} (-1)^{n-i} (x+i)^n = \sum_{i=0}^n \binom{n}{i} (-1)^{n-i} (x' + y + i)^n,$$

where $x' = x - y$. For y^n we can write

$$n! = \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} (y+k)^n,$$

By adding these two sums we get the equation for z^n .

$$2n! = \binom{n}{n} (-1)^0 z_{y+n}^n(x') - \binom{n}{n-1} (-1)^1 z_{y+n-1}^n(x') + \cdots + \binom{n}{0} (-1)^n z_y^n(x') \quad (2)$$

where $z_p^n(x') = (x' + p)^n + p^n$, $p \in \mathbb{N}$.

Notice that the coefficient before $n!$ is equal two in (2), it shows the behaviour of the function $z_p(x')$ and this can be useful. For $x' \geq 1$ obviously that

$$x' + p < z_p(x') < \sqrt[n]{2}(x' + p), \quad p \geq 1,$$

hence

$$1 < z_p(x') - z_{p-1}(x') < \sqrt[n]{2}. \quad (3)$$

With the growing of p the function $2(x' + p)^n$ grows more than the function $(x' + p)^n + p^n$. Also it is not hard to prove that $z_1(x') - z_0(x') < \sqrt[n]{2}$. Hereby, if we start from integer $z_0(x') = x'$ then $z_p(x')$ can be integer at least when

$$p(\sqrt[n]{2} - 1) > 1,$$

$$p > \frac{1}{\sqrt[n]{2} - 1}.$$

Therefore we can put the following statements for Fermat's Last Theorem:

Statement 1. *Fermat's Last Theorem is right when*

$$y < \frac{1}{\sqrt[n]{2} - 1}, \quad (x > y).$$

Proof. It has been proven before for every p hence it is proven for y .

□

Statement 2. *If the equation*

$$z_p^n(x') = (x' + p)^n + p^n, \quad (4)$$

where $x', p \geq 1$, has two integer roots $z_k(x')$ and $z_{k+i}(x')$ then

$$i > \frac{1}{\sqrt[n]{2} - 1}.$$

Proof. Suppose that $z_k(x')$ and $z_{k+i}(x')$ are integer roots of (4), then by (3)

$$i < z_{k+i}(x') - z_k(x') < \sqrt[n]{2} i,$$

hence

$$z_{k+i}(x') - z_k(x') - i < (\sqrt[n]{2} - 1)i,$$

but $z_{k+i}(x') - z_k(x') - i \geq 1$, therefore

$$i > \frac{1}{\sqrt[n]{2} - 1}.$$

□

The same we can say if $\{z_k(x')\}$ - the fractional part of $z_k(x')$ is less than $\sqrt[n]{2} - 1$. In this case the inequality for i takes the form

$$i > \frac{1}{\sqrt[n]{2} - 1} - 1.$$

This method is applicable to the whole kind of Diophantine equations. For example, lets take the following Diophantine equation

$$x^n + 1.15y^n = z^n,$$

where $n = 10^8$, $x, y \in \mathbb{N}$ and $x > y$. Applying the above method we can say that this equation has no integer roots for any x and if

$$y < \frac{1}{\sqrt[n]{2.15} - 1}.$$

If $x < y$ we can not apply this method directly because of the coefficient before y . In this case we have to start from $y' = 1$, $z_0(1) = \sqrt[n]{1.15}$ and then

$$z_p(1) = \sqrt[n]{1.15(1+p)^n + p^n}. \quad (5)$$

Hence in this case we have

$$y < \frac{2 - \sqrt[n]{1.15}}{\sqrt[n]{2.15} - 1}, \quad (x < y).$$

If we start from any y' in (5) then we have to start from the fractional part of $\sqrt[n]{1.15y'}$. In this case it depends on y' how many steps can be done. Let $\alpha = \{\sqrt[n]{1.15y'}\}$, $0 < \alpha < 1$, then

$$x < \frac{1 - \alpha}{\sqrt[n]{2.15} - 1}.$$

References

- [1] Brualdi R.: Introductory Combinatorics. China Machine Press, (2009).
- [2] Kleiner I.: From Fermat to Wiles: Fermat's Last Theorem Becomes a Theorem. Elem. Math., **55** (2000).
- [3] Wiles A.: Modular elliptic curves and Fermat's Last Theorem. Annals of Mathematics, **142** (1995), 443-551.

E-mail address: `khantarzhiiev@gmail.com`

MOSCOW, RUSSIA